



ParamectinTM

Pour-On for Cattle



Active Constituents

Abamectin 10 mg/mL

Actions

Paramectin Pour-On for Cattle contains abamectin, a member of the avermectin family of compounds which have a unique mode of action unrelated to that of other products used to control parasites of cattle in New Zealand. It provides a broad spectrum of efficacy against sensitive species of internal and external parasites of cattle. Resistance may develop to any drench.

Indications

For the treatment and control of internal and external parasites of cattle.

Paramectin Pour-On for Cattle, when used at the recommended dose level of 500 µg abamectin per kg liveweight, effectively controls the following parasites of cattle:

Gastrointestinal roundworms: adult and immature *Ostertagia* spp. (including inhibited immatures), *Haemonchus* spp., *Trichostrongylus* spp., *Cooperia* spp., *Bunostomum* spp., *Oesophagostomum* spp., and adults of *Chabertia* spp., *Nematodirus* spp., *Trichuris* spp. and *Strongyloides* spp.

Lungworms: adult and immature *Dictyocaulus viviparus*.

Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

Withholding Periods

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 35 days of last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.



Dosage and Administration

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose the mob to the heaviest animal by live weight in each group (bulls, cows, steers, calves etc.). Do not underdose. If there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Apply along the backline from shoulders to tail.

DOSE: 1 mL per 20 kg bodyweight (500 micrograms abamectin per kg bodyweight).

Liveweight	Dose	250 mL treats	500 mL treats	1 L treats	2.5 L treats	5 L treats	15 L treats
50 - 100 kg	5 mL	50 head	100 head	200 head	500 head	1000 head	3000 head
101 - 150 kg	7.5 mL	33 head	66 head	133 head	333 head	666 head	2000 head
151 - 200 kg	10 mL	25 head	50 head	100 head	250 head	500 head	1500 head
201 - 250 kg	12.5 mL	20 head	40 head	80 head	200 head	400 head	1200 head
251 - 300 kg	15 mL	16 head	33 head	66 head	166 head	333 head	1000 head
301 - 350 kg	17.5 mL	14 head	28 head	57 head	142 head	285 head	857 head
351 - 400 kg	20 mL	12 head	25 head	50 head	125 head	250 head	750 head
401 - 450 kg	22.5 mL	11 head	22 head	44 head	111 head	222 head	666 head
451 - 500 kg	25 mL	10 head	20 head	40 head	100 head	200 head	600 head
501 - 550 kg	27.5 mL	9 head	18 head	36 head	90 head	181 head	545 head
551 - 600 kg	30 mL	8 head	16 head	33 head	83 head	166 head	500 head
601 - 650 kg	32.5	7 head	15 head	30 head	76 head	153 head	461 head
Cattle in excess of 650 kg to be dosed at 1 mL per 20 kg bodyweight.							

Safety Directions

Warnings: : Harmful if swallowed. Causes serious eye irritation. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

Handling precautions: Obtain special instructions before use and do not handle until all safety precautions have been read and understood. Wear protective gloves/clothing and eye/face protection. Avoid contact during pregnancy and while nursing. Do not breathe vapours. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Wash contaminated clothing and/or protective equipment before reuse. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Collect spillage.



First Aid

For advice contact the National Poisons Centre on 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand. IF SWALLOWED: rinse mouth. Call a POISON CENTRE, or doctor if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. If eye irritation persists: Get medical advice. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice.

Presentation

Liquid: 5L, 15L

Storage

Store below 30°C (room temperature).

Classification

Non-restricted

Registration Number

ACVM No. A7726

EPA Approval Number

HSR001958