



Uro

Active Constituents

Prednil

Prednisolone 5 mg/tablet, chlorpheniramine maleate 2 mg/tablet.

Actions

Tablets

Combined anti-inflammatory, gluconeogenic and other metabolic regulatory effects of prednisolone and antihistamine properties of chlorpheniramine maleate.

Indications

For the treatment of allergic reactions and dermatitis and as an aid in the treatment of arthritic and rheumatic joint diseases in dogs and cats.

Contraindications

Corneal ulcers, diabetes mellitus, tuberculosis and renal insufficiency.

Precautions

Observe the usual precautions applicable to glucocorticoid therapy. Following prolonged glucocorticoid therapy, adrenocortical activity may be suppressed. It is therefore desirable to reduce treatment by lowering dose rates and frequency of administration.

Wound healing and fracture repair may be delayed, particularly in older animals.

Dosage and Administration

Dogs and cats: 1 tablet per 5 kg bodyweight daily, given orally.





First Aid

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone New Zealand 0800 764 766.

GHS Information

For GHS Information see Safety Data Sheet

Presentation

Tablets: 1,000s.

Storage

Store below 30°C (room temperature).

Classification

Restricted Veterinary Medicine (RVM). Only available under veterinary authorisation.

Registration Number

AVCM No. A003053



Products that work. People who care.

