



TECHNICAL NOTES

Strategik Combo

Dual Action Mineralised Sheep and Lamb Drench



Active Constituents

24.0 g/L Albendazole

37.5 g/L Levamisole HCl

0.25 g/L Cobalt

2.1 g/L Copper

0.55 g/L Zinc

0.5 g/L **Selenium**

Actions

For the treatment and control in sheep and lambs of benzimidazole or levamisole resistant roundworms. Also controls lungworms, tapeworms and adult liver fluke. Reduces the output of worm and fluke eggs.

STRATEGIK COMBO contains both albendazole, a member of the benzimidazole family, and levamisole. It is effective against internal parasites sensitive to one or both of these families. Resistance can develop to any anthelmintic. Correct drenching technique should be used. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.



TECHNICAL NOTES



Indications

STRATEGIK COMBO Dual Action Mineralised Sheep and Lamb Drench is intended for use against benzimidazole and levamisole susceptible worms of the following species:

ROUNDWORMS (mature and immature)

| | |
|-----------------------------|------------------------------------|
| Barber's pole worm | <i>Haemonchus contortus</i> |
| Small brown stomach worm | <i>Ostertagia</i> spp. |
| Stomach hair worm | <i>Trichostrongylus axei</i> |
| Black scour worm | <i>Trichostrongylus</i> spp. |
| Small intestinal worm | <i>Cooperia</i> spp. |
| Hookworm | <i>Bunostomum</i> spp. |
| Nodule worm | <i>Oesophagostomum columbianum</i> |
| Thin-necked intestinal worm | <i>Nematodirus</i> spp. |
| Large bowel worm | <i>Oesophagostomum venulosum</i> |
| Large mouthed bowel worm | <i>Chabertia ovina</i> |

LUNGWORMS

| | |
|----------------|-----------------------------|
| Large lungworm | <i>Dictyocaulus filaria</i> |
|----------------|-----------------------------|

TAPEWORMS

| | |
|--------------------|----------------------|
| Heads and Segments | <i>Moniezia</i> spp. |
|--------------------|----------------------|

| | |
|----------------------|--------------------------|
| LIVER FLUKE (Adults) | <i>Fasciola hepatica</i> |
|----------------------|--------------------------|

OVICIDAL

Reduces the output of viable worm and fluke eggs.

Precautions

Do not dose more frequently than 3-weekly. Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian.

Caution: In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper-poisoning (deaths) occurring following copper supplementation in this species.

Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.

Withholding Periods

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of cessation of the last treatment. Milk intended for sale for human consumption must be discarded during treatment and for not less than 10 days following the last treatment.



TECHNICAL NOTES

Dosage and Administration

By law, the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

General Directions:

Shake container well before use.

Administer orally, and check accuracy of drenching equipment before and during use. Clean drenching equipment thoroughly before and after use.

A representative sample of animals should be weighed before treatment.

Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Do not under dose.

Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Dose Rates

Sheep and Lambs: 1 mL per 5 kg body weight
(4.75 mg/kg albendazole, 7.5 mg/kg levamisole HCl).

Dose Volume Table

| Body Weight (kg) | Dose Volume (mL) | 1L pack treats (head) | 5L pack treats (head) | 20L pack treats (head) | 60L pack treats (head) |
|------------------|------------------|-----------------------|-----------------------|------------------------|------------------------|
| 15 | 3 | 333 | 1666 | 6666 | 20,000 |
| 20 | 4 | 250 | 1250 | 5000 | 15,000 |
| 25 | 5 | 200 | 1000 | 4000 | 12,000 |
| 30 | 6 | 166 | 833 | 3333 | 10,000 |
| 35 | 7 | 142 | 714 | 2857 | 8571 |
| 40 | 8 | 125 | 625 | 2500 | 7500 |
| 45 | 9 | 111 | 555 | 2222 | 6666 |
| 50 | 10 | 100 | 500 | 2000 | 6000 |
| 55 | 11 | 90 | 454 | 1818 | 5454 |
| 60 | 12 | 83 | 416 | 1666 | 5000 |
| 65 | 13 | 76 | 384 | 1538 | 4615 |
| 70 | 14 | 71 | 357 | 1428 | 4285 |
| 75 | 15 | 66 | 333 | 1333 | 4000 |

Sheep in excess of 75 kg to be dosed at the appropriate rate shown above the Dose Volume Table.





TECHNICAL NOTES



Safety Directions

Warnings: May be harmful if swallowed. Causes mild skin irritation. May cause an allergic skin reaction. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure.

Handling precautions: If medical advice is needed, have product container or label at hand. Obtain special instructions before use and do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing and eye/face protection. Do not breathe fumes/vapours. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse.

First Aid

If exposed, concerned or if you feel unwell: Call a Poisons Centre on 0800 POISON (0800 764 766) or a doctor. IF ON SKIN: wash with plenty of water. If skin irritation or rash occurs, get medical advice/attention.

Presentation

Liquid: 20 L

Storage

Store locked up, in the original container, tightly closed in a safe place. Store below 30°C (room temperature).

Registration Numbers

ACVM No. A7700