



## TECHNICAL NOTES

# Strategik Combo Plus Tape

Mineralised Combination Drench for Sheep



### Active Constituents

- 25.0 g/L Albendazole
- 37.5 g/L Levamisole HCl
- 18.8 g/L Praziquantel
- 0.2 g/L Cobalt (as disodium cobalt ethylenediamine tetra-acetate)
- 2.0 g/L Copper (as disodium copper ethylenediamine tetra-acetate)
- 0.6 g/L Zinc (as disodium zinc ethylenediamine tetra-acetate)
- 1.0 g/L Iodine (as potassium iodate)
- 0.5 g/L **Selenium** (as sodium selenate)

### Actions

A combination drench for the treatment and control in sheep and lambs of benzimidazole or levamisole resistant roundworms, tapeworms, lungworms and adult fluke.

STRATEGIK Combo plus Tape Mineralised Combination Drench for Sheep is a unique combination of praziquantel, a highly effective tapeworm agent, together with members of both the levamisole and benzimidazole families.

Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.





## TECHNICAL NOTES



### Indications

STRATEGIK Combo plus Tape Mineralised Combination Drench for Sheep is highly effective against the following parasites in sheep:

- **Heads and segments of tapeworms**
- **Mature and immature gastrointestinal roundworms:** *Haemonchus*, *Ostertagia*, *Trichostrongylus*, *Nematodirus*, *Cooperia*, *Strongyloides*, *Bunostomum*, *Oesophagostomum*, *Chabertia*, *Trichuris*.
- **Lungworms:** *Dictyocaulus*
- **Adult fluke**

### Precautions

Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian.

**Caution:** In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (death) occurring following copper supplementation in this species.

Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate liveweights carefully.

Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.

### Withholding Periods

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the food notice: Maximum Residue Levels for Agricultural Compounds.**

**Meat:** Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 10 days of cessation of the last treatment.

**Milk:** Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

### Dosage and Administration

**By law, the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

#### SHAKE WELL BEFORE USE

STRATEGIK Combo plus Tape Mineralised Combination Drench for Sheep is ready to use through standard drenching equipment. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. If there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Correct drenching technique should be used. Check the accuracy of drench guns regularly. Rinse gun after use.



## TECHNICAL NOTES

### Dose Rate

Sheep: 1 mL per 5 kg body weight

(7.5 mg/kg levamisole HCl, 5.0 mg/kg albendazole, 3.76 mg/kg praziquantel, 0.1 mg/kg **selenium**, 0.4 mg/kg copper, 0.12 mg/kg zinc, 0.04 mg/kg cobalt and 0.2 mg/kg iodine).

**Dose Volume Table**

Body Weight (kg)	Dose Volume (mL)	1 L treats (head)	5 L treats (head)	10 L treats (head)	20 L treats (head)
16 - 20	4	250	1250	2500	5000
21 - 25	5	200	1000	2000	4000
26 - 30	6	166	833	1666	3333
31 - 35	7	142	714	1428	2857
36 - 40	8	125	625	1250	2500
41 - 45	9	111	555	1111	2222
46 - 50	10	100	500	1000	2000
51 - 55	11	90	454	909	1818
56 - 60	12	83	416	833	1666
61 - 65	13	76	384	769	1538
66 - 70	14	71	357	714	1428
71 - 75	15	66	333	666	1333

Sheep in excess of 75 kg to be dosed at the appropriate rate shown above the Dose Volume Table.

### Safety Directions

**Warnings:** May be harmful if swallowed. May cause an allergic skin reaction. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure.

**Handling precautions:** Keep out of reach of children. Wear protective gloves/protective clothing/eye protection/face protection. Do not breathe vapours/spray. Contaminated clothing should not be allowed out of the workplace. Take off contaminated clothing and wash it before reuse.

### First Aid

IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention. IF exposed or concerned or if you feel unwell: Get medical advice/attention or contact the National Poisons Centre on 0800 POISON (0800 764 766). If medical advice is needed, have product container or label at hand.



## TECHNICAL NOTES

### Presentation

---

Liquid: 10 L

### Storage

---

Store in the original container, tightly closed, locked up in a safe place. Store below 30°C (room temperature).

### Registration Numbers

---

ACVM No. A9728

